



Accumulation of Organic Sunscreen Chemicals in Sharks and Teleost Fishes

Douglas H. Adams^{1*}, Ke He^{2*}, Ethan Hain², Jahir Antonio Batista Andrade², Anna Feerick², and Lee Blaney²

¹Florida Fish & Wildlife Conservation Commission, Fish & Wildlife Research Institute, Melbourne, FL

²University of Maryland Baltimore County, Baltimore, MD



Conclusions

- 1st analysis of sunscreen chemicals (UV-filters) in wild populations of Florida fishes;
- 7 UV-filters were detected in Bonnethead Shark muscle tissue from the Indian River Lagoon (IRL);
- 4-methylbenzylidene-camphor (4MBC) was present at high concentrations in Bonnetheads;
- 6 UV-filters were detected in IRL Red Drum tissue; and,
- Comparatively higher variation in UV filter distribution and concentrations among Red Drum individuals.

Background

- UV-filters have been widely used in sunscreens, pharmaceuticals, food packaging, and textiles;
- UV-filters are likely discharged into the IRL via wastewater effluent and recreational activity; and,
- Some UV-filters are endocrine-disrupting chemicals and can be toxic to corals, fish, and other aquatic organisms.

Objectives

- Build, refine, and optimize analytical techniques to measure a wide suite of UV-filters in estuarine fish muscle tissue; and,
- Analyze UV-filter concentrations in a wide array of fish species from the IRL and explore trophic level correlations.

Materials and Methods


- All fish samples were collected from the IRL by the FWC-FWRI's Fisheries-Independent Monitoring Program using 183-m haul seines;
- White dorsal muscle was collected using clean techniques;
- The tissue samples were shipped to UMBC on ice.




Bonnethead Shark, *Sphyrna tiburo*, collected from the IRL, FL





Red Drum, *Sciaenops ocellatus*, collected from the IRL, FL



- In total, 13 UV-filters were investigated in this study;
- Tissue samples were lyophilized and kept in -20 °C;
- Target analytes were extracted with a modified QuEChERS technique and cleaned by a novel reverse-SPE protocol;
- The concentration was determined by LC-MS/MS with mass-labeled surrogates and internal standards; and,
- Full details available in He *et al.*, 2017, *J. Chromatogr. A*




Modified QuEChERS extraction + reverse-SPE clean up




Liquid Chromatography tandem Mass Spectrometry (LC-MS/MS)

Key Findings



- 4MBC was present at high concentrations in Bonnetheads;
- In general, UV-filter concentrations in Red Drum tissue samples varied more than those in Bonnethead tissue, especially for BP3, BMDBM, and OS; and,
- Feeding ecology and trophic position likely play key roles in this variation.



Distribution and concentrations of UV filters in muscle from Bonnetheads (n=6, 608-913 mm PCL) and Red Drum (n=14, 322-542 mm SL). EHMC* is likely the metabolite of EHMC, but confirmation is needed.

- Bonnetheads have a comparatively restricted prey base (with blue crabs representing a high % of diet), where Red Drum have a more diverse diet and occupy a slightly lower trophic level within the IRL food web; and,
- In general, more hydrophobic UV-filters (e.g., HMS and BMDBM) exhibited higher concentrations in IRL fishes.

Future Research Directions

- Monitor UV-filters in water samples and identify potential sources;
- Expand sampling and analyses of UV-filters to other fish species representing all trophic levels within the IRL to further understand UV filter bioaccumulation within the ecosystem; and,
- Measure UV-filters in different fish organs to inform potential lethal and sub-lethal toxicity mechanisms

Acknowledgements

*Co-first authors. We greatly appreciate the efforts of scientist from FWC-FWRI's Fisheries-Independent Monitoring Program for assistance collecting samples for this study. This work was supported in part by funding from the U.S. Department of the Interior, U.S. Fish and Wildlife Service under Federal Aid in Sport Fish Restoration Project F-43, and from Florida's Saltwater Fishing Licenses. Analytical methods were developed using funding from the Maryland Sea Grant Program Development Fund.